

Source

Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4) is a chimeric monoclonal antibody recombinantly expressed from HEK293, which combines the variable region of a mouse monoclonal antibody with Human constant domain.

Clone

1B4

Isotype

Human IgG1 | Human Kappa

Conjugate

Unconjugated

Antibody Type

Recombinant Monoclonal

Reactivity

Virus

Immunogen

Recombinant Monkeypox virus (strain Zaire-96-I-16) A17L Protein is expressed from human 293 cells.

Specificity

Specifically recognizes Monkeypox virus (strain Zaire-96-I-16) A17L.

Application

| Application | Recommended Usage |
|-------------|-------------------|
| | |

ELISA 0.06-15 ng/mL

Purity

>90% as determined by SDS-PAGE.

Purification

Protein A purified/ Protein G purified

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE





Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4)


Catalog # A1L-M698



Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With Star Ribbon Pre-stained Protein Marker).

Bioactivity-ELISA


Monoclonal Anti-Monkeypox virus (strain Zaire-96-1-16) A17L Antibody, Human IgG1 (1B4) ELISA 0.1 µg of Monkeypox virus (strain Zaire-96-1-16) A17L Protein, His Tag per well



Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4) Conc. (ng/mL)

Immobilized Monkeypox virus (strain Zaire-96-I-16) A17L Protein, His Tag (Cat. No. A1L-M52H3) at 1 μ g/mL (100 μ L/well) can bind Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4) (Cat. No. A1L-M698) with a linear range of 0.06-2 ng/mL (QC tested).

Bioactivity-SPR



Monoclonal Anti-Monkeypox virus (strain Zaire-96-I-16) A17L Antibody, Human IgG1 (1B4) (Cat. No. A1L-M698) captured on Protein A Chip can bind Monkeypox virus (strain Zaire-96-I-16) A17L Protein, His Tag (Cat. No. A1L-M52H3) with an affinity constant of 7.98 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Monkeypox is a rare zoonosis caused by monkeypox virus, which has become the most serious orthpoxvirus and consists of complex double stranded DNA. The cases are mostly in central and western Africa. The pathogenesis of monkeypox is that the virus invades the body from respiratory mucosa, multiplies in lymphocytes, and incurs into blood producing transient venereal toxemia. after the virus multiplies in cells, the cells can invade the blood and propagate to the skin of the whole body, causing lesions. The envelope glycoprotein A35R on the EV surface has been predicted to influence intercellular diffusion of virions.

Clinical and Translational Updates

