

Mouse CD79B Protein, Fc Tag (MALS verified)

Catalog # CDB-M5253



BIOSYSTEMS
Acro

Synonym

CD79b,B29,IGB,Ig-beta

Source

Mouse CD79B, Fc Tag(CDB-M5253) is expressed from human 293 cells (HEK293). It contains AA Val 26 - Asp 158 (Accession # [P15530-1](#)).

Predicted N-terminus: Val 26

Molecular Characterization

CD79B(Val 26 - Asp 158) P15530-1	Fc(Pro 100 - Lys 330) P01857
-------------------------------------	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 40.3 kDa. The protein migrates as 33 kDa and 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse CD79B, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS



The purity of Mouse CD79B, Fc Tag (Cat. No. CDB-M5253) is more than 85% and the molecular weight of this protein is around 85-105 kDa verified by SEC-MALS.

[Report](#)

Background

B-cell antigen receptor complex-associated protein beta chain (CD79b) is also known as B-cell-specific glycoprotein B29, Ig-beta, Immunoglobulin-associated B29 protein, B29 and IGB, which is a single-pass type I membrane protein containing one Ig-like V-type (immunoglobulin-like) domain and one ITAM domain. CD79b is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR). CD79b can enhance

Discounts, Gifts,
and more!



Mouse CD79B Protein, Fc Tag (MALS verified)

Catalog # CDB-M5253



phosphorylation of CD79A, possibly by recruiting kinases which phosphorylate CD79A or by recruiting proteins which bind to CD79A and protect it from dephosphorylation. Defects in CD79b are the cause of agammaglobulinemia type 6 (AGM6) that is a primary immunodeficiency characterized by profoundly low or absent serum antibodies and low or absent circulating B cells due to an early block of B-cell development.

Clinical and Translational Updates

Discounts, Gifts,
and more!



www.acrobiosystems.com

3/21/2024