Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Monomer, MALS verified)

Catalog # HLA-H82E8





Source

Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI)
Complex Protein(HLA-H82E8) is expressed from human 293 cells (HEK293). It
contains AA Gly 25 - Ile 308 (HLA-A*02:01) & Ile 21 - Met 119 (B2M) &
FMVFLQTHI peptide (Accession # <u>AAA59606.1</u> (HLA-A*02:01) & <u>P61769-1</u>
(B2M) & FMVFLQTHI).

Predicted N-terminus: Gly 25 & Ile 21

Molecular Characterization

Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein is produced by co-expression of HLA and B2M loaded with EBV EBNA peptide.

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (AvitagTM).

The protein has a calculated MW of 36.3 kDa and 11.7 kDa. The protein migrates as 41-43 kDa and 10 kDa when calibrated against <u>Star Ribbon Prestained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using AvitagTM technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from $0.22~\mu m$ filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

SEC-MALS



The purity of Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) is more than 90% and the molecular weight of this protein is around 45-60 kDa verified by SEC-MALS.

Report



Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Monomer, MALS verified)


Catalog # HLA-H82E8





Bioactivity-ELISA


Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA 0.1 μ g of Anti-HLA class I Antibody, Human IgG1 (W6/32) per well



Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein Conc. (ng/mL)


Immobilized Anti-HLA class I Antibody, Human IgG1 (W6/32) at 1 μ g/mL (100 μ L/well) can bind Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) with a linear range of 0.5-8 ng/mL (QC tested).

Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA 0.1 μg of Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein per well



Immobilized Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) at 1 μ g/mL (100 μ L/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 μ g/well) plate can bind Anti-B2M Antibody, Human IgG1 with a linear range of 0.2-3 ng/mL (Routinely tested).

Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA 0.1 μg of Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein per well



Immobilized Biotinylated Human HLA-A*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) at 1 μg/mL (100 μL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 μg/well) plate

can bind Anti-HLA class I Antibody, Human IgG1 (W6/32) with a linear range

of 0.2-3 ng/mL (Routinely tested).

Clinical and Translational Updates

