

Synonym

LGR5, GPR49, GPR67, G-protein coupled receptor 49, G-protein coupled receptor 67, G-protein coupled receptor HG38, Leucine-rich repeat-containing G-protein coupled receptor 5

Source

Cynomolgus LGR5, His Tag(LG5-C52H7) is expressed from human 293 cells (HEK293). It contains AA Ser 22 - Gly 557 (Accession # [G7PI19-1](#)).
Predicted N-terminus: Ser 22

Molecular Characterization

LGR5(Ser 22 - Gly 557)
G7PI19-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus

The protein has a calculated MW of 61.7 kDa. The protein migrates as 66-70 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.



This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Cynomolgus LGR5, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Cynomolgus LGR5, His Tag (Cat. No. LG5-C52H7) at 5 µg/mL (100 µL/well) can bind Human R-Spondin 1 (21-146), Fc Tag (Cat. No. RS1-H5269) with a linear range of 0.2-5 ng/mL (QC tested).

Immobilized Human R-Spondin 1 (21-146), Fc Tag (Cat. No. RS1-H5269) at 5 µg/mL (100 µL/well) can bind Cynomolgus LGR5, His Tag (Cat. No. LG5-C52H7) with a linear range of 0.1-10 ng/mL (Routinely tested).

Background

LGR5 (also known as GPR49) is a seven-transmembrane protein of the class A Rhodopsin-like family of GPCRs. LGR5 and LGR4 bind the R-spondins with high affinity and mediate the potentiation of Wnt/ β -catenin signaling by enhancing Wnt-induced LRP6 phosphorylation. The LGR5/RSPO complex could promote Wnt signalling via the neutralization of two transmembrane E3 ligases, RNF43 and ZNRF3.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.