



### Synonym

Streptavidin,SA

### Source

Streptavidin Protein-Acrininium ester (STN-NA114) is Acrininium ester chemically conjugated Streptavidin expressed from E. coli cells.

### Molecular Characterization

Streptavidin carries no "tag". The protein has a calculated MW of 13.8 kDa. The protein migrates as 15-16 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE).

### Labeling

*Acrininium ester, can react with the primary amino group of protein. Under alkaline conditions, NHS is replaced as the leaving group, and the protein forms a stable amide bond with Acrininium ester.*

### Protein Ratio

Passed as determined by binding MPCLIA.

### Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please protect from light and avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


### SDS-PAGE



Streptavidin Protein-Acrininium ester on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

### Bioactivity-MPCLIA

### SEC-MALS



The purity of Streptavidin Protein-Acrininium ester (Cat. No. STN-NA114) is more than 90% and the molecular weight of this protein is around 55-75 kDa verified by SEC-MALS.


[Report](#)

Discounts, Gifts,  
and more!






**PD-1 binding with PD-L1 by MPCLIA**




Immobilized 0.04 µg/Test of Biotinylated Human PD-1 Protein, Avitag,His Tag (Cat. No. PD1-H82E4) to the Streptavidin Protein-Acrininium ester (Cat. No. STN-NA114, 0.008 µg/Test), incubated with 20 µL/Test of Human PD-L1 Protein, Mouse IgG1 Fc Tag (Cat. No. PD1-H52A3) at increasing concentration coupled to Anti-Mouse IgG Magnetic Beads (10 µg beads/Test). Detection was performed with sensitivity of 7.8 ng/mL in Magnetism particulate chemiluminescence immunoassay (MPCLIA) (KEYSMILE, SMART 6500S) (QC tested).

**Detection of Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM122) by MPCLIA**



Immobilized 0.04 µg/Test of Biotinylated SARS-CoV-2 Spike RBD, His,Avitag (Cat. No. SPD-C82E9) to the Streptavidin Protein-Acrininium ester (Cat. No. STN-NA114, 0.008 µg/Test), incubated with 20 µL/Test of Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM122) (Cat. No. S1N-M12A1) at increasing concentration coupled to Protein G Magnetic Beads (10 µg beads/Test). Detection was performed with sensitivity of 0.244 ng/mL in Magnetism particulate chemiluminescence immunoassay (MPCLIA) (KEYSMILE, SMART 6500S) (Routinely tested).

**Detection of Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM122) by MPCLIA**



Immobilized 0.04 µg/Test of Biotinylated SARS-CoV-2 Spike RBD, His,Avitag (Cat. No. SPD-C82E9) to the Streptavidin Protein-Acrininium ester (Cat. No. STN-NA114, 0.008 µg/Test), incubated with 20 µL/Test of Anti-SARS-CoV-2 Spike RBD Antibody, Chimeric mAb, Human IgG1 (AM122) (Cat. No. S1N-M12A1) at increasing concentration coupled to Anti-Human IgG Magnetic Beads (10 µg beads/Test). Detection was performed with sensitivity of 7.8 ng/mL in Magnetism particulate chemiluminescence immunoassay (MPCLIA) (KEYSMILE, SMART 6500S) (Routinely tested).

**Background**

Streptavidin is a tetrameric protein purified from the bacterium *Streptomyces avidinii*, and exhibits high binding affinity for biotin. Able to bind one molecule of biotin with each subunit. Streptavidin (PI=6.0-7.5) has lower level of non-specific binding to various biological components at physiological pH than avidin (PI=7.4), resulting from its isoelectric point (PI).Streptavidin is useful in affinity chromatography, ELISA, immunohistochemistry and Western Blotting.

**Clinical and Translational Updates**

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.

Discounts, Gifts, and more!

